



Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Submitted to Physical Review C

PION-CONDENSATION THRESHOLD IN NUCLEAR MATTER
AND THERMAL Δ -ISOBARS


P. Hecking

February 1981

RECEIVED
LAWRENCE
BERKELEY LABORATORY

APR 23 1981

LIBRARY AND
DOCUMENTS SECTION



DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Pion-Condensation Threshold in Nuclear Matter
and Thermal Δ -Isobars*

P. Hecking
Nuclear Science Division
Lawrence Berkeley Laboratory
University of California
Berkeley, CA 94720

Abstract: The pion-condensation threshold in symmetric ($N = Z$) nuclear matter at finite temperatures $0 \leq k_B T \leq 100$ MeV is calculated. The influence of thermally excited Δ -isobars--present in the nucleon medium--on the threshold is investigated and found to be moderately repulsive (i.e. raising the threshold density).

[Nuclear structure, pion-condensation in nuclear matter,
thermal Δ -isobars at finite temperature]

PACS-index: 21.65 + f, 24.90. + d

Since the pioneering work of Migdal¹⁾ much attention, both in theory and experiment, has been directed to exotic states of baryon matter, such as pion-condensation. Heavy-ion

*This work was supported by the Director, Office of Energy Research, Division of Nuclear Physics of the Office of High Energy and Nuclear Physics of the U.S. Department of Energy under Contract W-7405-ENG-48.

collisions at high bombarding energies promise the chance to observe this phenomenon on the earth, whereas conclusions about pion-condensed cores of neutron stars from their cooling are rather indirect and plagued by uncertainties about the equation of state, superfluidity and competing cooling mechanisms.²⁾

In previous calculations concerning pion-condensation, the Δ -isobar occurs either in a Δ -nucleon hole excitation by a virtual pion³⁾ or as an admixture to the nucleons, resulting from a full diagonalisation of the baryon system in the presence of a pion-condensate.^{4,5)} At finite temperature in nuclear matter, there are already purely thermal baryon excitations present, even in the absence of a pion-condensate. The most important of them, because of its lowest excitation energy, is the Δ -isobar. Consequently, at finite temperature the (negative) pion self-energy Π results in part from the interaction of a pion with a Fermi-sea of Δ -isobars, as indicated in fig. 1.

The threshold of pion-condensation is given by the minimal baryon density ρ_B , required for a solution of the equation

$$\omega^2 = k^2 + m_\pi^2 + \Pi(k, \omega, \rho_B, T) \quad (1)$$

with $\omega = 0$ in spin-isospin symmetric ($N = Z$) nuclear matter. The baryon density ρ_B is the sum of the nucleon density and the Δ -density

$$\rho_B = \rho_N + \rho_\Delta \quad (2)$$

with the Δ -density ρ_Δ given by

$$\rho_{\Delta} = \rho_B \cdot 4 \exp\left(-\frac{\omega_{\Delta}}{k_B T}\right) \left[1 + 4 \exp\left(-\frac{\omega_{\Delta}}{k_B T}\right)\right]^{-1} \quad (3)$$

where $\omega_{\Delta} = 2.3 m_{\pi}$ is the mass difference between nucleon and Δ -isobar. The factor 4 results from the four times larger degeneracy of the Δ relative to the nucleon. The pion self-energy consists of an s-wave and a p-wave part, the latter modified by short-range repulsive interactions, incorporated in the Fermi-liquid parameter g' :

$$\Pi = \Pi_s + \Pi_p \quad \Pi_p = \Pi_p^{(0)} [1 - g' k^{-2} \Pi_p^{(0)}]^{-1} \quad \Pi_s = 0.03 \rho \rho_0^{-1} m_{\pi}^2 \quad (4)$$

with the nuclear matter density $\rho_0 = 0.17 \text{ fm}^{-3}$

The zeroth order p-wave self-energy $\Pi_p^{(0)}$ is given by a sum of Lindhard-functions:

$$\Pi_p^{(0)} = -k^2 m_{\pi}^{-2} [f^2(k^2) U_N + f^{*2}(k^2) (U_{N\Delta} + U_{\Delta N}) + g^{*2}(k^2) U_{\Delta}] \quad (5)$$

For symmetric nuclear matter the nucleon Lindhard function U_N is given by the principal part integral

$$U_N = 4P \int d^3p (2\pi)^{-3} [n(p) - n(p+k)] \cdot [\epsilon(p+k) - \epsilon(p)]^{-1} \quad (6)$$

with the Fermi distribution

$$n(p, T) = [1 + \exp(\epsilon(p) - \mu(T))/k_B T]^{-1} \quad (7)$$

where $\epsilon(p) = p^2 (2m^*)^{-1}$ is the nucleon energy with an effective mass m^* . The chemical potential $\mu(T)$ is determined from the nucleon density

$$\rho_N = 4 \int d^3p (2\pi)^{-3} n(p, \mu, T) \quad (8)$$

Furthermore, the Δ -nucleon hole excitation function $U_{N\Delta}$ is given by

$$U_{N\Delta} = \frac{16}{9} \int d^3p (2\pi)^{-3} 2n(p) [\varepsilon_{\Delta}(p+k) - \varepsilon(p)]^{-1} \quad (9)$$

The Δ -single particle energy is $\varepsilon_{\Delta}(p) = p^2(2m_{\Delta})^{-1} + \omega_{\Delta}$, with $m_{\Delta} = 9.01 m_{\pi}$ and $\omega_{\Delta} = 2.3 m_{\pi}$. The functions $U_{\Delta N}$ and U_{Δ} have the same structure as their counterparts $U_{N\Delta}$ and U_N ; only the roles of nucleon and Δ -isobar are exchanged.

This means the following replacements in the above functions:

$n \rightarrow n_{\Delta}$	$m^* \rightarrow m_{\Delta}^*$	
$\mu \rightarrow \mu_{\Delta}$	$m_{\Delta} \rightarrow m^*$	(10)
$(\mu_{\Delta} \text{ determined from } \rho_{\Delta})$	$\omega_{\Delta} \rightarrow -\omega_{\Delta}$	

The spin-isospin factor 16/9 in $U_{N\Delta}$ is the same for $U_{\Delta N}$, whereas that for U_{Δ} is 25 instead of 4 for U_N . In eq. (5), the finite range vertex structure is represented by form factors:

$$\frac{f(k^2)}{f} = \frac{f^*(k^2)}{f^*} = \frac{g^*(k^2)}{g^*} = \frac{\Lambda^2 - m^2}{\Lambda^2 + k^2} \quad (11)$$

with the coupling constants $f^2 = 4\pi \cdot 0.08$ and $f^* = 2f$ and $g^* = 4/5f$ according to the Chew-Low-model.⁶⁾

In earlier calculations^{3,5)} it turned out already that the critical density for pion-condensation is sensitive to the effective nucleon mass m^* , the short-range correlation parameter g' , the vertex cutoff Λ , and the temperature $k_B T$. Measurements and calculations of these quantities are mostly concentrated at nuclear matter density ρ_0 with very meager information about the density region above. The cutoff Λ is

rather well determined in the range 1-1.4 GeV⁷⁾ and presumably not very sensitive to the density; we adopt the two values 1.2 and 1.4 GeV for our model calculation. The Fermi-liquid parameter g' and the effective mass m^* are not well known at density $\rho > \rho_0$. At $\rho = \rho_0$, g' has values of 0.6-0.7⁸⁾. In spite of a few model calculations up to $\rho \sim 2\rho_0$, its value remains uncertain above ρ_0 . We chose the three values 0.5, 0.6 and 0.7 in order to cope with this uncertainty. For the effective mass m^* , which is probably of all three quantities most sensitive to the density, we make two choices: a density-independent $m^* = 0.8 m_N$, which is characteristic for ρ_0 and a density dependent $m^*(\rho) = m_N(1 - k_F(\rho)k_F(\rho_0)^{-1} 0.2)$, which gives $m^* = 0.8 m_N$ at ρ_0 and is an average of several possibilities, offered in ref. 9). The quantity k_F is the Fermi momentum.

The results for the critical baryon density ρ_c are given in fig. 2. At zero temperature, where no Δ -isobars are present, the results are in good agreement with earlier calculations^{3,4,5)}. One obvious feature is the rather "parallel" behavior of the phase boundaries with increasing temperature for various cut-offs Λ and parameters g' . Although both quantities have much influence on the critical density, this influence is rather constant between $0 \leq k_B T \leq 100$ MeV. This is not the case for the two different m^* . At low temperature, $m^*(\rho) < 0.8 m_N$ gives a smaller pion self-energy and consequently a larger ρ_c . Above $k_B T \sim 50$ MeV another effect gains importance: the "softening"

of the Fermi surface by the temperature is relatively smaller for $m^*(\rho)$, because it gives a larger Fermi energy $k_F^2(2m^*(\rho))^{-1}$ than $m^* = 0.8 m_N$. Both effects tend to cancel with the result that both choices of m^* give nearly identical phase boundaries for $k_B T \gtrsim 50$ MeV.

The Fermi-sea of thermally excited Δ -isobars is of importance only above $k_B T \sim 50$ MeV. Below this temperature, the Δ -density approaches zero exponentially. In eq. (5) the Lindhard-function U_Δ is attractive, whereas $U_{\Delta N}$ is repulsive. The depopulation of the nucleon Fermi-sea in favor of the Δ -isobar is also repulsive, all three together resulting in an increase of the critical density, which is not very dramatic, however; it is a less than 20% effect, even at $k_B T \sim 100$ MeV. The temperature 100 MeV is reached in a typical central heavy-ion collision with bombarding energy ~ 1 GeV/N, provided complete thermalization is obtained.

We conclude that a Fermi-sea of thermally excited Δ -isobars increases the critical density for pion-condensation in nuclear matter but is below $k_B T \sim 100$ MeV of less importance than the vertex cutoff Λ and the parameter g' .

Of all five considered quantities, Δ -admixture, effective nucleon mass m^* , temperature $k_B T$, vertex cutoff Λ , and short-range correlation parameter g' , the last one still remains the most decisive and unfortunately also (especially at $\rho > \rho_0$) the least well known.

This work was supported in part by the DFG under contract He 1155/1 and by the the Director, Office of Energy Research, Division of Nuclear Physics of the Office of High Energy and Nuclear Physics of the U.S. Department of Energy under Contract W-7405-ENG-48.

References

1. A.B. Migdal, Nucl. Phys. A210, 421 (1973); Rev. Mod. Phys. 50, 107 (1978)
2. O. Maxwell, Astrophys. J. 231, 201 (1979)
3. V. Ruck, M. Gyulassy and W. Greiner, Z. Phys. A277, 391 (1976)
4. H.J. Pirner, M. Rho, K. Yazaki and P. Bonche, Nucl. Phys. A329, 491 (1979)
5. P. Hecking, Nucl. Phys. A348, 493 (1980)
6. G.E. Brown and W. Weise, Phys. Rep. 22C, 279 (1975)
7. J.W. Durso, A.D. Jackson and B.J. Verwest, Nucl. Phys. A282, 404 (1977)
8. S.O. Bäckman, O. Sjöberg and A.D. Jackson, Nucl. Phys. A321, 10 (1979)
E. Oset and M. Rho, Phys. Rev. Lett. 42, 47 (1979)
J. Meyer-ter-Vehn, PLT 34, Max Planck-Gesellschaft, Munich 1980
9. Y. Futami, H. Toki and W. Weise, Phys. Lett. 77B, 37 (1978)


Figure Captions

Fig. 1. Lowest order p-wave pion self-energy $\Pi_p^{(0)}$.



(a) Direct nucleon Born term and Δ -nucleon hole excitation. (b) Direct Δ Born term and nucleon- Δ hole excitation. Crossed terms omitted for simplicity.

Fig. 2. Critical density $\rho_c(T)$ as function of the temperature and various parameters. The pion-condensed phase is to the right of the curves. Without Δ -Fermi-sea: full curves: $\Lambda = 1.2$ GeV and $m^* = 0.8$; dashed curves: $\Lambda = 1.4$ GeV and $m^* = 0.8$; dotted curves: $\Lambda = 1.2$ GeV and $m^* = m(\rho)$. With Δ -Fermi-sea: dash-dotted curves: $\Lambda = 1.2$ GeV and $m^* = 0.8$. Short-range correlation parameter g' as indicated in the figure.

XBL812-350



b



a





Fig. 1



XBL812-349

Fig. 2